

PAIN MANAGEMENT



GUNA®-THORACIC is a homeopathic complex medicine in injectable ampoules designed for infiltration treatment of acute and chronic pain of different origin, affecting the posterior area of the spinal column.

USES

Thoracic pain due to cartilage degenerative thoracic spine disorders (thoracic osteoarthritis in association with GUNA®-POLYARTHRITIS), Thoracic pain due to scoliosis, Thoracic pain due to thoracic long muscle trigger points (in association with GUNA®-MUSCLE), Pain due to thoracic spine osteophytosis, Simple posterior thoracic pain (thoracic syndrome), Mechanical imbalance (costo-vertebral facet joint syndrome), Thoracic spinal ligament syndrome, Thoracic spinal nerve root pain (in association with GUNA®- NEURAL), Pain from spinal osteoporosis.

GUNA®-THORACIC may be used together with other homeopathic medications.

INGREDIENTS

Active ingredients:

Chininum sulphuricum 8X HPUS 4 parts; Anti interleukin 1 alpha 4C, Anti interleukin 1 beta 4C, Beta-Endorphin 4C, Calcarea phosphorica 6X HPUS, Cimicifuga racemosa 4X HPUS, Intervertebral disk, Porcine 4X, Spongia tosta 10X HPUS 2 parts; Chelidonium majus 4X HPUS, Conium maculatum 6X HPUS 1 part.

Inactive ingredient:

Sterile isotonic sodium chloride solution.

Ingredients that are not included in the HPUS are manufactured according to the European Pharmacopeia and the Official German Homeopathic Pharmacopeia.

DOSAGE FORMS

2 ml glass ampoules in package of 10 ampoules
2 ml glass ampoules in package of 50 ampoules

DIRECTIONS

STANDARD PROTOCOL FOR I.M. ADMINISTRATION

1 ampoule 1-3 times a week according to severity and clinical evolution (1-2 months of treatment).

STANDARD PROTOCOL ACCORDING TO MESOTHERAPEUTIC TECHNIQUE

Using 1-2 ampoules per treatment: 2 treatments for the first 2 weeks, 1 treatment a week till pain relief (average 8-10 sessions). For chronic pathologies: continue 1 treatment a week for 1 month till pain relief, then 1 treatment a month.

Select application site according to trigger points, tender points, referred pain zones, acupuncture points, nerve key points, Head zones or "local pain points".

METHOD OF ADMINISTRATION

In case of subcutaneous administration, use a 13 mm, 30G, a 4 mm, 27G needle or an insulin needle to apply the classic mesotherapeutic technique.

Administration may vary according to individual needs.